

COVID-19 and kids

How does COVID-19 affect kids?



All children are at risk of getting COVID-19, but the good news is their symptoms are generally mild. Many children experience a cough, fever, and a runny nose, and only require rest at home, recovering quickly.

A very small percentage of children experience a barking cough, prolonged fever, breathing difficulties and abdominal pain, and these children are advised to see their doctor.




In Australia in 2021, **less than 3%** of children aged under 16 years who had a COVID-19 infection were hospitalised due to COVID-19.¹ Many were admitted because their parents were too unwell to look after them.²

Children with underlying health conditions are at higher risk of experiencing severe COVID-19 symptoms. Conditions include asthma, obesity, prematurity, and compromised immune systems.²⁻⁵ Children aged younger than 6 months are also at greater risk.^{2,3}

What about Omicron?

Is this strain a bigger problem for kids?

Most children who get Omicron experience a mild infection and early data on hospitalisation rates shows **less than 2% of children aged under 10 years who get Omicron are hospitalised.**⁶ Early data indicates a lower risk of hospitalisation among Omicron cases in school-aged children compared to the Delta strain.^{6,7}



Are there any long-term impacts?

Long-term impacts of COVID-19 are still being investigated but early research suggests that **persistent symptoms can occur for many weeks in a small group of children.**⁸ These include the following symptoms:



Headache



Fatigue



Sleep disturbance



Concentration difficulties



Abdominal pain

About 1 in 2,500 children with COVID-19 can develop a condition called **Multisystem Inflammatory Disease in Children (MIS-C)** after having COVID-19.² The majority of children who are hospitalised with MIS-C are not vaccinated.^{9,10}

How can COVID-19 be prevented?

COVID-19 is spread by breathing in air or small droplets exhaled by an infected person, or touching your eyes, nose or mouth with hands that have COVID-19 germs on them. **It's important to teach children ways how to reduce the risk of contracting COVID-19, such as:**



Physical distancing



Frequent hand washing with soap, or using hand sanitiser



Wearing well-fitted masks over mouth and nose
(Link to mask fact sheet)



Staying home when unwell



Covering coughs and sneezes



COVID-19 vaccination

If my child gets COVID-19, how do I care for them?

If your child becomes unwell with COVID-19, you can provide pain relief such as **paracetamol**, especially in the case of fever, ensure they **drink plenty of fluids** and encourage them to **rest** until they have recovered.¹¹

If symptoms gets worse (e.g., fever for 5+ days, breathing difficulties, abdominal pain, drinking less, sleepy, irritable), **seek medical attention**.

You should also try and minimise the spread of COVID-19 in your home by **washing hands regularly** and wiping down commonly touched surfaces, such as benches, taps and door handles with a disinfectant or soap and water. **Thoroughly wash any common household items used by the person with COVID-19**, such as cups, eating utensils, bedding, and towels, and consider using separate bathroom facilities if available.¹¹



For more information on
COVID-19 in kids visit
tacklingcovid19.org.au

References

1. COVID-19 National Incident Room Surveillance Team. COVID-19 Australia: Epidemiology Report 55. Communicable Diseases Intelligence. 2021;45
2. Williams P, Koirala A, Saravanos G, et al. COVID-19 in children in NSW, Australia, during the 2021 Delta outbreak: severity and disease spectrum. medRxiv. 2022.
3. Hobbs CV, Woodworth K, Young CC, et al. Frequency, characteristics and complications of COVID-19 in hospitalized Infants. The Pediatric Infectious Disease Journal. 2021.
4. Mukkada S, Bhakta N, Chantada GL, et al. Global characteristics and outcomes of SARS-CoV-2 infection in children and adolescents with cancer (GRCCC): a cohort study. The Lancet Oncology. 2021;22(10):1416-1426.
5. Haeusler GM, Ammann RA, Carlesse F, et al. SARS-CoV-2 in children with cancer or after haematopoietic stem cell transplant: An analysis of 131 patients. European Journal of Cancer. 2021;159:78-86
6. NSW Health. COVID-19 weekly surveillance in NSW: epidemiological week 52 ending 1 January 2022. 2022 [cited 2022 Jan]. Available from: <https://www.health.nsw.gov.au/Infectious/covid-19/Documents/covid-19-surveillance-report-20220113.pdf>
7. UK Health Security Agency. SARS-CoV-2 variants of concern and variants under investigation in England. Technical briefing: Update on hospitalisation and vaccine effectiveness for Omicron VOC-21NOV-01 (B.1.1.529) 2021 [cited 2022 Jan]. Available from: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1045619/Technical-Briefing-31-Dec-2021-Omicron_severity_update.pdf
8. Zimmermann P, Pittet LF, Curtis N. How common is long COVID in children and adolescents? The Pediatric Infectious Disease Journal. 2021;40(12):e482
9. Zambrano LD, Newhams MM, Olson SM, et al. Effectiveness of BNT162b2 (Pfizer-BioNTech) mRNA vaccination against multisystem inflammatory syndrome in children among persons aged 12–18 Years—United States, July–December 2021. 2022
10. Levy M, Recher M, Hubert H, et al. Multisystem inflammatory syndrome in children by COVID-19 vaccination status of adolescents in France. JAMA. 2021
11. Health Direct. Caring for people with COVID-19: Australian Government Department of Health; 2021 [cited 2022 Jan]. Available from: <https://www.healthdirect.gov.au/coronavirus-covid-19-information-for-carers-faqs>
12. Attwell K, Carlson SJ, Tchilingirian J, Harper T, McKenzie L, Roberts L, Rizzi M, Westphal D, Effler P, Hughes C, Swift V, Blyth CC. Coronavax: Preparing Community and Government for COVID-19 Vaccination: a Research Protocol for a mixed-methods social research project. BMJ Open. 2021;11(6):e049356